

The variational principle of topological pressure for actions of sofic groupoids

Xiaoyao Zhou^{,†} Ercai Chen^{†‡}

[†] *School of Mathematical Sciences and Institute of Mathematics, Nanjing Normal University,*

Nanjing 210023, P.R.China, [‡] Center of Nonlinear Science,

Nanjing University, Nanjing 210093, P.R.China.

e-mail: zhouxiaoyaodeyouxian@126.com

ecchen@nju.edu.cn

Abstract. This article establishes the variational principle of topological pressure for actions of sofic groupoids.

Keywords and phrases Topological pressure; sofic groupoid; variational principle.

1 Introduction

Variational principles are beautiful results in a dynamical system. Establishing variational principle is an important topic in dynamical system theory. The first variational principle that reveals the relationship between topological entropy and measure-theoretic entropy was obtained by L. Goodwyn [8] and T. Goodman [7]. M. Misiurewicz gave a short proof in [17]. R. Bowen [4] established the variational principle of entropy for non-compact set in 1973. Y. Pesin and B. Pitskel [19] studied the variational principle of pressure for non-compact set.

Romagnoli [20] introduced two types of measure-theoretic entropies relative to a finite open cover and proved the variational principle between local entropy and one type. Later, Glasner and Weiss [6] proved that if the system is invertible, then the local variational principle is also true for another type measure-theoretic entropy. W. Huang & Y. Yi [11] obtained the variational principle of local pressure. W. Huang, X. Ye and G. Zhang [10] established the variational principle of local entropy for a

countable discrete amenable group action. B. Liang and K. Yan [16] generalized it to local pressure for sub-additive potentials of amenable group actions.

Recently, L. Bowen [1], [2] introduced a notion of entropy for measure-preserving actions of a countable discrete sofic group on a standard probability space admitting a generating finite partition. Just after that, D. Kerr and H. Li [13] established variational principle of entropy for sofic group actions. G. Zhang [23] generalized it to the variational principle of local entropy for countable infinite sofic group actions. N. Chung [5] generalized the result of D. Kerr and H. Li [13] to the variational principle of pressure for sofic group actions.

Very recently, L. Bowen [3] generalized the sofic entropy theory to class-bijective extensions of sofic groupoids, and established the variational principle between topological entropy and measure entropy.

This article establishes the variational principle of topological pressure for actions of sofic groupoids.

2 Preliminaries

The section presents some basic definitions and notations about groupoid.

2.1 Discrete groupoid

A groupoid $\mathcal{H}^1$ is a set consists of morphisms, equipped with a set of objects $\mathcal{H}^0$, source and range maps $\varsigma, \tau : \mathcal{H}^1 \rightarrow \mathcal{H}^0$, an injective inclusion map $i : \mathcal{H}^0 \rightarrow \mathcal{H}^1$, such that

- (i) for any $x \in \mathcal{H}^0$, $\varsigma(i(x)) = \tau(i(x)) = x$;
- (ii) let $\mathcal{H}^2 = \{(f, g) \in \mathcal{H}^1 \times \mathcal{H}^1 : \varsigma(f) = \tau(g)\}$ and define a composition map $c : \mathcal{H}^2 \rightarrow \mathcal{H}^1$ satisfying
- (iii) for any $(f, g) \in \mathcal{H}^2$, $\varsigma(c(f, g)) = \varsigma(g)$, $\tau(c(f, g)) = \tau(f)$;
- (iv) for any $f \in \mathcal{H}^1$, there exists a unique element $f^{-1} \in \mathcal{H}^1$ with $c(f^{-1}, f) = i(\varsigma(f))$ and $c(f, f^{-1}) = i(\tau(f))$.

For convenience, a remark about the notations of a groupoid to be used in this article is presented here.

Remark 2.1. Let $\mathcal{H}^1$ be a groupoid defined as above.

- Write $\mathcal{H}$ instead of $\mathcal{H}^1$, fg instead of $c(fg)$ for the sake of convenience.
- Let $[[\mathcal{H}]]$ be the collection of all Borel subsets $f \subset \mathcal{H}$, whose source and range maps restricted to f are Borel isomorphisms onto their respective images.
- Given $f, g \in [[\mathcal{H}]]$, set $f^{-1} = \{h^{-1} : h \in f\}$ and $fg = \{h \in \mathcal{H} : h = f'g' \text{ for some } f' \in f, g' \in g\}$.

- $\mathcal{H} \in [[\mathcal{H}]]$ and any Borel subset P of $\mathcal{H}^0$ belongs to $[[\mathcal{H}]]$ (Using the inclusion map, $\mathcal{H}^0$ can be viewed as a subset of $\mathcal{H}$.).
- If $f \in [[\mathcal{H}]]$ and $x \in \varsigma(f)$, then fx is well-defined, and $fx = \varsigma^{-1}(x) \cap f$. However, fx may not be contained in $\mathcal{H}^0$. To avoid this, we let $f \cdot x = \tau(fx) \in \mathcal{H}^0$. In a similar way, if $P \subset \mathcal{H}^0$, then we let $f \cdot P = \tau(fP)$.
- Let $[\mathcal{H}] \subset [[\mathcal{H}]]$ be the collection of all Borel subsets $f \subset \mathcal{H}$, whose source and range maps are each Borel isomorphisms onto $\mathcal{H}^0$. Composition makes $[[\mathcal{H}]]$ an inverse semi-group called the semi-group of partial automorphisms, $[\mathcal{H}]$ a group called the full group of $\mathcal{H}$.
- Let $[\mathcal{H}]_{top}$ be the collection of all closed sets $f \subset \mathcal{H}$ such that the restrictions of the source and range maps to f are homeomorphisms onto $\mathcal{H}^0$. Under composition, $[\mathcal{H}]_{top}$ is a subgroup of $[\mathcal{H}]$.

A measurable groupoid $\mathcal{H}$ means a groupoid $\mathcal{H}$ is equipped with the structure of a standard Borel space satisfying that $\mathcal{H}^0$ is a Borel set and the source, range, composition and inversion maps are all Borel.

A groupoid $\mathcal{H}$ is discrete means that for any $x \in \mathcal{H}^0$, $\varsigma^{-1}(x)$ and $\tau^{-1}(x)$ are countable.

A discrete probability measure groupoid refers to a discrete measurable groupoid $\mathcal{H}$ together with a Borel probability measure ν over $\mathcal{H}^0$ satisfying that if ν_ς, ν_τ are the measures over $\mathcal{H}$ defined by

$$\nu_\varsigma(B) = \int_{\mathcal{H}^0} |\varsigma^{-1}(x) \cap B| d\nu(x), \nu_\tau(B) = \int_{\mathcal{H}^0} |\tau^{-1}(x) \cap B| d\nu(x),$$

for any Borel set $B \subset \mathcal{H}$, then ν_ς is equivalent to ν_τ . In particular, if $\nu_\varsigma = \nu_\tau$, then $(\mathcal{H}, \nu)$ is pmp (probability-measure-preserving). The article deals with pmp groupoids. Hence, we let ν denote $\nu_\varsigma = \nu_\tau$, and ν restricted to $\mathcal{H}^0$ is ν .

A discrete topological groupoid is a discrete groupoid $\mathcal{H}$ paired with a topology such that the structure maps (source, range, inverse and composition) are continuous. An open subset $f \subset \mathcal{H}$ is called a bisection means that the restrictions of the source and range maps to f are homeomorphisms onto their images which are open subsets of $\mathcal{H}^0$. $\mathcal{H}$ is étale refers to that if every $g \in \mathcal{H}$ is contained in a bisection.

Remark 2.2. Assume that $(\mathcal{H}, \nu)$ is pmp.

- Given a Borel set $A \subset \mathcal{H}^0$, set $\partial A = A \cap \overline{\mathcal{H}^0 \setminus A}$.
- Let $\mathcal{B}_\partial(\mathcal{H}^0, \nu)$ be the set of all Borel subset $A \subset \mathcal{H}^0$ satisfying $\nu(\partial A) = \nu(\partial(\mathcal{H}^0 \setminus A)) = 0$.

- Let $[[\mathcal{H}]]_{top}$ be the collection of all elements of $[[\mathcal{H}]]$ with the form $f = \cup_{i=1}^n f_i$, where
 - (i) for each i there is a bisection U_i with $f_i \subset U_i$,
 - (ii) $\{\varsigma(f_i)\}_{i=1}^n \subset \mathcal{B}_\partial(\mathcal{H}^0, \nu)$ are pairwise disjoint,
 - (iii) $\{\tau(f_i)\}_{i=1}^n \subset \mathcal{B}_\partial(\mathcal{H}^0, \nu)$ are pairwise disjoint.
- For $f \in [[\mathcal{H}]]$, the trace of f is given by $tr_{\mathcal{H}}(f) = \nu(\mathcal{H}^0 \cap f)$. Also $|f|_{\mathcal{H}} = \nu(f)$.

Example [3] Let $d \in \mathbb{N}$. The full groupoid on $\{1, \dots, d\}$ is $\Delta_d := \{1, \dots, d\}^2$ and the unit space is $\Delta_d^0 := \{(i, i) : 1 \leq i \leq d\}$. The structure maps are given by $\varsigma(i, j) = (j, j)$, $\tau(i, j) = (i, i)$, $(i, j)^{-1} = (j, i)$ and $(i, j)(j, k) = (i, k)$. Let $\zeta_d(E) = |E|/d$ for any set $E \subset \Delta_d$. (Δ_d, ζ_d) is a pmp groupoid. We simplify $[\Delta_d]$ as $[d]$ that is isomorphic with the symmetric group on $\{1, \dots, d\}$ and simplify $[[\Delta_d]]$ as $[[d]]$ that is the set of all subsets $f \subset \Delta_d$ such that $\varsigma : f \rightarrow \varsigma(f)$ and $\tau : f \rightarrow \tau(f)$ are bijections. Let $tr_d := tr_{\Delta_d}$, $|\cdot|_d := |\cdot|_{\Delta_d}$ and $tr_d(f) = |f \cap \Delta_d^0|/d$ and $|f|_d = |f|/d$.

2.2 Sofic approximations, actions, spanning and separating sets

Let $(\mathcal{H}, \nu)$ denote a pmp discrete groupoid. Given $d > 0$, let $\text{Map}([[\mathcal{H}]], [[d]])$ be the collection of all Borel maps from $[[\mathcal{H}]]$ to $[[d]]$. For a finite set $F \subset [[\mathcal{H}]]$ and $\sigma : [[\mathcal{H}]] \rightarrow [[d]]$, set

$$N(\sigma, F) = \{\sigma' \in \text{Map}([[\mathcal{H}]], [[d]]) : \sigma'(f) = \sigma(f), \forall f \in F\}.$$

The Borel structure of $\text{Map}([[\mathcal{H}]], [[d]])$ can be viewed as generated by all such $N(\sigma, F)$.

$X \subset_f Y$ means that X is a finite subset of Y . Let $F \subset_f [[\mathcal{H}]]$, $\delta > 0$, a map $\sigma : [[\mathcal{H}]] \rightarrow [[d]]$ is (F, δ) -multiplicative means for any $s, t \in F$, $|\sigma(st)\Delta\sigma(s)\sigma(t)|_d < \delta$ and (F, δ) -trace-preserving refers to $|tr_d(\sigma(s)) - tr_{\mathcal{H}}(s)| < \delta$ for any $s \in F$.

Definition 2.1. [3] (*Sofic approximation*) Let J be a direct set. For each $j \in J$, let $d_j \in \mathbb{N}$ and $\mathbb{P}_j$ be a Borel probability measure on $\text{Map}([[\mathcal{H}]], [[d_j]])$. The family $\mathbb{P} = \{\mathbb{P}_j\}_{j \in J}$ is a sofic approximation to $(\mathcal{H}, \nu)$ means

(i) for any $F \subset_f [[\mathcal{H}]]$ and $\delta > 0$,

$$\lim_{j \rightarrow J} \mathbb{P}(\{\sigma \in \text{Map}([[\mathcal{H}]], [[d_j]]) : \sigma \text{ is } (F, \delta) \text{ - trace - preserving}\}) = 1.$$

(ii) for any $F \subset_f [[\mathcal{H}]]$, $\delta > 0$, there exists $j \in J$ such that $j' \geq j$ implies $\mathbb{P}_{j'}$ -almost every σ is (F, δ) -multiplicative.

(iii) $\lim_{j \rightarrow J} d_j = +\infty$.

The groupoid $(\mathcal{H}, \nu)$ is sofic means it has a sofic approximation.

Let $\mathcal{G}, \mathcal{H}$ be measurable groupoid. A map $\pi : \mathcal{G} \rightarrow \mathcal{H}$ is a groupoid morphism means that for every $(f, g) \in \mathcal{G}^2$, $\pi(fg) = \pi(f)\pi(g)$, for each $f \in \mathcal{G}$, $\pi(f)^{-1} = \pi(f^{-1})$ and $\pi(\mathcal{G}^0) \subset \mathcal{H}^0$. It is class-bijective if for every $a \in \mathcal{G}^0$, the restriction of π to $\varsigma^{-1}(a)$ is a bijection onto $\varsigma^{-1}(\pi(a))$ and the restriction of π to $\tau^{-1}(a)$ is also a bijection onto $\tau^{-1}(\pi(a))$. If π is also surjective, then $\mathcal{G}$ is a class-bijective extension of $\mathcal{H}$ or, equivalently, $\mathcal{H}$ is a class-bijective factor of $\mathcal{G}$. π is pmp (probability-measure-preserving) if $\pi_*\mu = \nu$ and $(\mathcal{G}, \mu), (\mathcal{H}, \nu)$ are pmp groupoid.

If $\pi : (\mathcal{G}, \mu) \rightarrow (\mathcal{H}, \nu)$ is class-bijective then $\pi^{-1} : [[\mathcal{H}]] \rightarrow [[\mathcal{G}]]$ is a homeomorphism. Assume $\mathcal{G}$ and $\mathcal{H}$ are topological groupoid, π is continuous and pmp then $\pi^{-1}([[\mathcal{H}]]_{top}) \subset [[\mathcal{G}]]_{top}$. Given $x \in \mathcal{G}$ and $f \in [[\mathcal{H}]]$, $fx := \pi^{-1}(f)x$. If $x \in \mathcal{G}^0$ then we let $f \cdot x$ be $\tau(\pi^{-1}(f)x) = \tau(fx)$.

Definition 2.2. Given a pseudo-metric space (Z, ρ) and $\epsilon > 0$, a subset $Y \subset Z$ is (ρ, ϵ) -separated means that for any $y_1 \neq y_2 \in Y$, $\rho(y_1, y_2) > \epsilon$. Given $X \subset Z$, let $N_\epsilon(X, \rho)$ be the maximum cardinality of a (ρ, ϵ) -separated subset $Y \subset X$. For $X, Y \subset Z$, $Y(\rho, \epsilon)$ -spans X means that for each $x \in X$, there exists $y \in Y$ with $\rho(x, y) < \epsilon$. Let $N'_\epsilon(X, \rho)$ be the minimum cardinality of a set $Y \subset Z$ which (ρ, ϵ) -spans X .

3 Topological pressure

Assume $\mathcal{G}, \mathcal{H}$ are two discrete separable topological groupoid such that $\mathcal{G}^0$ and $\mathcal{H}^0$ are compact metrizable spaces, $\pi : \mathcal{G} \rightarrow \mathcal{H}$ is a continuous class-bijective factor map, $(\mathcal{H}, \nu)$ is a pmp groupoid. Given a sofic approximation $\mathbb{P} = \{\mathbb{P}_j\}_{j \in J}$ to $(\mathcal{H}, \nu)$, $\varphi \in C(\mathcal{G}^0, \mathbb{R})$, a bias β and $p \in [1, \infty]$, we define the sofic topological pressure of π with respect to $(\mathbb{P}, p, \beta, \varphi)$, via a choice of generating pseudo-metric.

For an integer $d > 0$, we write $x \in (\mathcal{G}^0)^d$ as $x = (x_1, \dots, x_d)$. For $f \in [[\mathcal{H}]]$, we set $f \cdot x := \{f \cdot x_1, \dots, f \cdot x_d\}$. $f \cdot x_i$ is not defined when $\pi(x_i) \notin \varsigma(f)$. When this occurs, we set $f \cdot x_i := \star$, where $\star$ is a special symbol. So $f \cdot x \in (\mathcal{G}^0 \cup \{\star\})^d$.

Δ_d^0 is viewed as $\{1, \dots, d\}$ and given $\sigma : [[\mathcal{H}]] \rightarrow [[d]]$, $f \in [[\mathcal{H}]]$ and $i \in \{1, \dots, d\}$, we write $\sigma(f)i \in \{1, \dots, d\}$ instead of $\sigma(f) \cdot i$. With x as above, we define $x \circ \sigma(f) := (x_{\sigma(f)1}, \dots, x_{\sigma(f)d})$. If $i \notin \varsigma(\sigma(f))$, then $x_{\sigma(f)i}$ is not well-defined. When it occurs, we let $x_{\sigma(f)i} := \star$. Thus $x \circ \sigma(f) \in (\mathcal{G}^0 \cup \{\star\})^d$.

Choose a continuous pseudo-metric ρ on $\mathcal{G}^0$. ρ can be extended to $\mathcal{G}^0 \cup \{\star\}$ by setting $\rho(\star, \star) = 0$ and $\rho(\star, x) = \max\{\rho(y, z) : y, z \in \mathcal{G}^0\}$ for any $x \in \mathcal{G}^0$. Two pseudo-metrics on $(\mathcal{G}^0 \cup \{\star\})^d$ are given by

$$\rho_2(x, x') := \left(\frac{1}{d} \sum_{i=1}^d \rho(x_i, x'_i)^2 \right)^{1/2}, \quad \rho_\infty(x, x') := \max_{1 \leq i \leq d} \rho(x_i, x'_i).$$

Definition 3.1. [3] (*Approximate partial orbits*) Let $C(\mathcal{H}^0)$ be the space of continuous complex-valued functions on $\mathcal{H}^0$. For a map $\sigma : [[\mathcal{H}]] \rightarrow [[d]]$, finite sets $F \subset [[\mathcal{H}]]$, $K \subset C(\mathcal{H}^0)$ and $\delta > 0$, set

$$\text{Orb}_\nu(\pi, \sigma, F, K, \delta, \rho) := \{(x_1, \dots, x_d) \in (\mathcal{G}^0)^d : \delta > \rho_2(f \cdot x, x \circ \sigma(f)), \forall f \in F, \\ \delta > |d^{-1} \sum_{i=1}^d k(\pi(x_i)) - \int k d\nu|, \forall k \in K\}.$$

Definition 3.2. [3] A bias β for J is either an element of $\{-, +\}$ or a nonprincipal ultrafilter on J . Given a function $\Phi : J \rightarrow \mathbb{R}$, if β is an ultrafilter then the ultralimit $\lim_{j \rightarrow \beta} \Phi(j)$ is well-defined. Otherwise, define

$$\lim_{j \rightarrow \beta} \Phi(j) := \begin{cases} \liminf_{j \rightarrow J} \Phi(j), & \text{if } \beta = -; \\ \limsup_{j \rightarrow J} \Phi(j), & \text{if } \beta = +. \end{cases}$$

Let

$$M_\epsilon(\text{Orb}_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_2, \varphi) := \sup_{\mathcal{E}} \left\{ \sum_{(x_1, \dots, x_d) \in \mathcal{E}} \exp\left(\sum_{i=1}^d \varphi(x_i)\right) \right\},$$

where $\mathcal{E}$ runs over (ρ_2, ϵ) -separated subsets of $\text{Orb}_\nu(\pi, \sigma, F, K, \delta, \rho)$.

If $1 \leq p < \infty$, then we define

$$\|M_\epsilon(\text{Orb}_\nu(\pi, \cdot, F, K, \delta, \rho), \rho_2, \varphi)\|_{p, \mathbb{P}_j} = \left(\int M_\epsilon(\text{Orb}_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_2, \varphi)^p d\mathbb{P}_j(\sigma) \right)^{1/p}.$$

Define topological pressure by separated set as follows:

$$P_{\mathbb{P}, p}^\beta(\pi, \rho, 2, \varphi) := \sup_{\epsilon > 0} \inf_{\delta > 0} \inf_{F \subset_f [[\mathcal{H}]]_{\text{top}}} \inf_{K \subset_f C(\mathcal{H}^0)} \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|M_\epsilon(\text{Orb}_\nu(\pi, \cdot, F, K, \delta, \rho), \rho_2, \varphi)\|_{p, \mathbb{P}_j}, \\ P_{\mathbb{P}, p}^\beta(\pi, \rho, \infty, \varphi) := \sup_{\epsilon > 0} \inf_{\delta > 0} \inf_{F \subset_f [[\mathcal{H}]]_{\text{top}}} \inf_{K \subset_f C(\mathcal{H}^0)} \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|M_\epsilon(\text{Orb}_\nu(\pi, \cdot, F, K, \delta, \rho), \rho_\infty, \varphi)\|_{p, \mathbb{P}_j}.$$

Remark 3.1. We are suppressing the choice of bias β and parameter $p \in [1, \infty]$ from the notation. Thus, $P_{\mathbb{P}, p}^\beta(\pi, \rho, 2, \varphi)$ can be simplified as $P_{\mathbb{P}}(\pi, \rho, 2, \varphi)$ and $P_{\mathbb{P}, p}^\beta(\pi, \rho, \infty, \varphi)$ can be expressed by $P_{\mathbb{P}}(\pi, \rho, \infty, \varphi)$.

Remark 3.2. The quantity $\frac{1}{d_j} \log \|M_\epsilon(\text{Orb}_\nu(\pi, \cdot, F, K, \delta, \rho), \rho_2, \varphi)\|_{p, \mathbb{P}_j}$ is monotone increasing in δ and monotone decreasing in ϵ, F, K . This implies that the supremum and the infimums can be replaced by the appropriate limits.

Let

$$M'_\epsilon(\text{Orb}_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_2, \varphi) := \inf_{\mathcal{E}'} \left\{ \sum_{(x_1, \dots, x_d) \in \mathcal{E}'} \exp\left(\sum_{i=1}^d \varphi(x_i)\right) \right\},$$

where $\mathcal{E}'$ runs over (ρ_2, ϵ) -spanning subsets of $\text{Orb}_\nu(\pi, \sigma, F, K, \delta, \rho)$.

Lemma 3.1. *Topological pressure defined by spanning set is equal to given by separated set. More precisely,*

$$P_{\mathbb{P}}(\pi, \rho, 2, \varphi) = \sup_{\epsilon > 0} \inf_{\delta > 0} \inf_{F \subset_f [[\mathcal{H}]]_{top}} \inf_{K \subset_f C(\mathcal{H}^0)} \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|M'_\epsilon(Orb_\nu(\pi, \cdot, F, K, \delta, \rho), \rho_2, \varphi)\|_{p, \mathbb{P}_j}, (*)$$

$$P_{\mathbb{P}}(\pi, \rho, \infty, \varphi) = \sup_{\epsilon > 0} \inf_{\delta > 0} \inf_{F \subset_f [[\mathcal{H}]]_{top}} \inf_{K \subset_f C(\mathcal{H}^0)} \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|M'_\epsilon(Orb_\nu(\pi, \cdot, F, K, \delta, \rho), \rho_\infty, \varphi)\|_{p, \mathbb{P}_j} (**).$$

Proof. The proof of $(*)$ is presented here. Since $(**)$ is similar to $(*)$, the proof is omitted.

Let $\mathcal{E} \subset Orb_\nu(\pi, \sigma, F, K, \delta, \rho)$ be a (ρ_2, ϵ) -separated subset with maximal cardinality, then $\mathcal{E}(\rho_2, 2\epsilon)$ -spans $Orb_\nu(\pi, \sigma, F, K, \delta, \rho)$. Therefore,

$$\begin{aligned} M_\epsilon(Orb_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_2, \varphi) \\ &\geq \sum_{(x_1, \dots, x_d) \in \mathcal{E}} \exp\left(\sum_{i=1}^d \varphi(x_i)\right) \\ &\geq M'_{2\epsilon}(Orb_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_2, \varphi). \end{aligned}$$

This implies that

$$P_{\mathbb{P}}(\pi, \rho, 2, \varphi) \geq \sup_{\epsilon > 0} \inf_{\delta > 0} \inf_{F \subset_f [[\mathcal{H}]]_{top}} \inf_{K \subset_f C(\mathcal{H}^0)} \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|M'_\epsilon(Orb_\nu(\pi, \cdot, F, K, \delta, \rho), \rho_2, \varphi)\|_{p, \mathbb{P}_j}.$$

Next, the proof of the opposite inequality is presented as follows. For any $x', y' \in \mathcal{G}^0$, there exists $\eta_1 > 0$ such that $|\varphi(x') - \varphi(y')| < \eta_1$, whenever $\rho(x', y') < \sqrt{\epsilon}$. There exists (ρ_2, ϵ) -separated subset $\mathcal{E}_1 \subset Orb_\nu(\pi, \sigma, F, K, \delta, \rho)$ such that

$$M_\epsilon(Orb_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_2, \varphi) \leq \sum_{(x_1^1, \dots, x_d^1) \in \mathcal{E}_1} \exp\left(\sum_{i=1}^d \varphi(x_i^1)\right) \cdot \exp(1)$$

and there exists a $(\rho, \epsilon/2)$ -spanning subset $\mathcal{E}_2$ of $Orb_\nu(\pi, \sigma, F, K, \delta, \rho)$ such that

$$M'_{\epsilon/2}(Orb_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_2, \varphi) \cdot \exp(1) \geq \sum_{(x_1^2, \dots, x_d^2) \in \mathcal{E}_2} \exp\left(\sum_{i=1}^d \varphi(x_i^2)\right).$$

Define $\phi : \mathcal{E}_1 \rightarrow \mathcal{E}_2$ by choosing, for each $(x_1^1, \dots, x_d^1) \in \mathcal{E}_1$, some point $\phi((x_1^1, \dots, x_d^1)) = (x_1^2, \dots, x_d^2)$ with $\rho_2((x_1^1, \dots, x_d^1), (x_1^2, \dots, x_d^2)) < \epsilon/2$. Note that ϕ is injective and $\rho_2((x_1^1, \dots, x_d^1), (x_1^2, \dots, x_d^2)) < \epsilon/2$ implies that $|\{1 \leq i \leq d : \rho(x_i^1, x_i^2) \geq \sqrt{\epsilon}\}| \leq \lceil \frac{d\epsilon}{4} \rceil$. Therefore,

$$\begin{aligned}
& M'_{\epsilon/2}(Orb_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_2, \varphi) \cdot \exp(1) \\
& \geq \sum_{(x_1^2, \dots, x_d^2) \in \mathcal{E}_2} \exp\left(\sum_{i=1}^d \varphi(x_i^2)\right) \\
& \geq \sum_{(x_1^2, \dots, x_d^2) \in \phi\mathcal{E}_1} \exp\left(\sum_{i=1}^d \varphi(x_i^2)\right) \\
& = \sum_{\substack{(x_1^1, \dots, x_d^1) \in \mathcal{E}_1 \\ (x_1^2, \dots, x_d^2) = \phi((x_1^1, \dots, x_d^1))}} \exp\left(\sum_{i=1}^d \varphi(x_i^1) + \sum_{i=1}^d \varphi(x_i^2) - \sum_{i=1}^d \varphi(x_i^1)\right) \\
& \geq \sum_{(x_1^1, \dots, x_d^1) \in \mathcal{E}_1} \exp\left(\sum_{i=1}^d \varphi(x_i^1)\right) \exp\left(-(d - \lceil \frac{d\epsilon}{4} \rceil)\eta_1 - 2\|\varphi\| \lceil \frac{d\epsilon}{4} \rceil\right) \\
& \geq M_\epsilon(Orb_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_2, \varphi) \exp\left(-(d - \lceil \frac{d\epsilon}{4} \rceil)\eta_1 - 2\|\varphi\| \lceil \frac{d\epsilon}{4} \rceil - 1\right).
\end{aligned}$$

This implies

$$P_{\mathbb{P}}(\pi, \rho, 2, \varphi) \leq \sup_{\epsilon > 0} \inf_{\delta > 0} \inf_{F \subset_f [[\mathcal{H}]]_{top}} \inf_{K \subset_f C(\mathcal{H}^0)} \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|M'_\epsilon(Orb_\nu(\pi, \cdot, F, K, \delta, \rho), \rho_2, \varphi)\|_{p, \mathbb{P}_j}.$$

□

Remark 3.3. In the definitions of $M_\epsilon(Orb_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_2, \varphi)$ it suffices to take the supremum over those (ρ_2, ϵ) separated sets with maximal cardinality. Similarly, in the definitions of $M'_\epsilon(Orb_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_2, \varphi)$, it suffices to take the infimum over the (ρ_2, ϵ) spanning sets with minimal cardinality.

Lemma 3.2. $P_{\mathbb{P}}(\pi, \rho, 2, \varphi) = P_{\mathbb{P}}(\pi, \rho, \infty, \varphi)$.

Proof. Since any (ρ_2, ϵ) -separated set is also (ρ_∞, ϵ) -separated, we get

$$M_\epsilon(Orb_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_2, \varphi) \leq M_\epsilon(Orb_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_\infty, \varphi),$$

for any σ, F, K, δ . This implies that $P_{\mathbb{P}}(\pi, \rho, 2, \varphi) \leq P_{\mathbb{P}}(\pi, \rho, \infty, \varphi)$.

Next, we prove the opposite direction. Let $\frac{1}{10} > \epsilon > 0, \Omega$ be a $(\rho, \sqrt{\epsilon})$ -spanning subset of $\mathcal{G}^0$ with minimum cardinality. Let $\delta > 0, F \subset_f [[\mathcal{H}]], K \subset_f C(\mathcal{H}^0), \sigma : [[\mathcal{H}]] \rightarrow [[d]]$ and $\mathcal{E} \subset (\mathcal{G}^0)^d$ be a (ρ_2, ϵ) -spanning set for $Orb_\nu(\pi, \sigma, F, K, \delta, \rho)$ with

$$2M'_\epsilon(Orb_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_2, \varphi) \geq \sum_{(y_1, \dots, y_d) \in \mathcal{E}} \exp\left(\sum_{i=1}^d \varphi(y_i)\right).$$

Let $\eta = \lceil \epsilon d \rceil$. Define $\mathcal{E}' \subset (\mathcal{G}^0)^d$ as follows:

For $y = (y_1, \dots, y_d) \in \mathcal{E}$, every set $\Lambda \subset [d]$ of cardinality η and every map $\phi : \Lambda \rightarrow \Omega$, y^ϕ is given by

$$y_i^\phi = \begin{cases} y_i, & i \notin \Lambda; \\ \phi(i), & i \in \Lambda. \end{cases}$$

Let $\mathcal{E}'$ be the collection of all y^ϕ over all such $y \in \mathcal{E}$ and $\phi : \Lambda \rightarrow \Omega$. Then

$$\sum_{y^\phi \in \mathcal{E}'} \exp\left(\sum_{i=1}^d \varphi(x_i)\right) \leq \binom{d}{\eta} |\Omega|^\eta \sum_{y \in \mathcal{E}} \exp\left(\sum_{i=1}^d \varphi(y_i) + 2\|\varphi\|\eta\right).$$

We claim that $\mathcal{E}'$ is $(\rho_\infty, \sqrt{\epsilon})$ -spanning for $Orb_\nu(\pi, \sigma, F, K, \delta, \rho)$. In fact, let $z \in Orb_\nu(\pi, \sigma, F, K, \delta, \rho)$. Since $\mathcal{E}$ is (ρ_2, ϵ) -spanning for $Orb_\nu(\pi, \sigma, F, K, \delta, \rho)$, there exists $y \in \mathcal{E}$ such that $\rho_2(y, z) \leq \epsilon$, that is,

$$\frac{1}{d} \sum_{i=1}^d \rho(y_i, z_i)^2 \leq \epsilon^2.$$

So, there exists a set $\Lambda \subset [d]$ such that for $i \notin \Lambda$, $\rho(y_i, z_i) \leq \sqrt{\epsilon}$ and $|\Lambda| = \eta$. By the definition of Ω , for every $i \in \Lambda$, there exists a point $\phi(i) \in \Omega$ such that $\rho(\phi(i), z_i) \leq \sqrt{\epsilon}$. Thus, $\rho_\infty(y^\phi, z) \leq \sqrt{\epsilon}$, that shows that $\mathcal{E}'$ is $(\rho_\infty, \sqrt{\epsilon})$ -spanning for $Orb_\nu(\pi, \sigma, F, K, \delta, \rho)$. It follows that

$$\begin{aligned} & M'_\epsilon(Orb_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_\infty, \varphi) \\ & \leq \sum_{y^\phi \in \mathcal{E}'} \exp\left(\sum_{i=1}^d \varphi(y_i^\phi)\right) \\ & \leq \binom{d}{\eta} |\Omega|^\eta \sum_{y \in \mathcal{E}} \exp\left(\sum_{i=1}^d \varphi(y_i) + 2\|\varphi\|\eta\right) \\ & \leq 2 \exp(2\|\varphi\|\eta) \binom{d}{\eta} |\Omega|^\eta \cdot M'_\epsilon(Orb_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_2, \varphi). \end{aligned}$$

Then the desired result follows from Stirling's approximation formula. $\square$

Definition 3.3. [3] A pseudo-metric ρ on $\mathcal{G}^0$ is dynamically generating for $\pi : \mathcal{G} \rightarrow \mathcal{H}$ means that for each $x, y \in \mathcal{G}^0$ there exists $f \in [\mathcal{H}]_{top}$ with $\rho(f \cdot x, f \cdot y) > 0$.

Theorem 3.1. Let ρ, ρ' be two dynamically generating continuous pseudo-metrics on $\mathcal{G}^0$. Then $P_{\mathbb{P}}(\pi, \rho, \varphi) = P_{\mathbb{P}}(\pi, \rho', \varphi)$.

Remark 3.4. The proof of Theorem 3.1 uses only properties (ii) and (iii) of the definition of sofif approximation.

Definition 3.4. Given Theorem 3.1, the sofic topological pressure of π with respect to $(\mathbb{P}, p, \beta, \varphi)$ is given by $P_{\mathbb{P}}(\pi, \varphi) := P_{\mathbb{P}}(\pi, \rho, \varphi) = P_{\mathbb{P}, p}^{\beta}(\pi, \varphi)$, where ρ is any dynamically generating continuous pseudo-metric on $\mathcal{G}^0$.

It is worth mentioning that this is the relative pressure with respect to the measure ν . Because the sofic approximation $\mathbb{P}$ determines ν , ν is implicitly referenced in the notation.

Lemma 3.3. If ρ, ρ' are two continuous metrics on $\mathcal{G}^0$, then $P_{\mathbb{P}}(\pi, \rho, \varphi) = P_{\mathbb{P}}(\pi, \rho', \varphi)$.

Proof. Because ρ and ρ' are continuous metrics on $\mathcal{G}^0$, and $\mathcal{G}^0$ is compact, for every $\delta > 0$ and sufficiently large integer $n \gg 0$, there exist $\delta_0, \epsilon_n > 0$ such that

- $\rho'(x, y) \leq \sqrt{\delta_0} \Rightarrow \rho(x, y) \leq \delta$,
- $\delta_0 \leq \delta^2$;
- $\rho'(x, y) \geq \epsilon_n \Rightarrow \rho(x, y) \geq \frac{1}{n}$;
- $\lim_{n \rightarrow \infty} \epsilon_n = 0$.

Let $\Omega = \max\{\rho(x, y) : x, y \in \mathcal{G}^0\}$ be the diameter of ρ . We claim that for any $\sigma : [[\mathcal{H}]] \rightarrow [[d]]$, $F \subset_f [[\mathcal{H}]]_{top}$ and $K \subset_f C(\mathcal{H}^0)$,

$$Orb_{\nu}(\pi, \sigma, F, K, \delta_0, \rho') \subset Orb_{\nu}(\pi, \sigma, F, K, \delta(\Omega^2 + 1)^{1/2}, \rho),$$

the proof of which is presented in Lemma 6.9 of [3]. By choice of ϵ_n ,

$$M_{\epsilon_n}(Orb_{\nu}(\pi, \sigma, F, K, \delta_0, \rho'), \rho'_{\infty}, \varphi) \leq M_{\frac{1}{n}}(Orb_{\nu}(\pi, \sigma, F, K, \delta(\Omega^2 + 1)^{1/2}, \rho), \rho_{\infty}, \varphi).$$

Thus,

$$\begin{aligned} & \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|M_{\epsilon_n}(Orb_{\nu}(\pi, \cdot, F, K, \delta_0, \rho'), \rho'_{\infty}, \varphi)\|_{p, \mathbb{P}_j} \\ & \leq \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|M_{\frac{1}{n}}(Orb_{\nu}(\pi, \cdot, F, K, \delta(\Omega^2 + 1)^{1/2}, \rho), \rho_{\infty}, \varphi)\|_{p, \mathbb{P}_j}. \end{aligned}$$

Taking the infimum over $\delta_0 > 0$ then over all $\delta > 0$, then over all $F \subset_f [[\mathcal{H}]]_{top}$ and $K \subset_f C(\mathcal{H}^0)$, then the supremum over all n , we have $P_{\mathbb{P}}(\pi, \rho', \varphi) \leq P_{\mathbb{P}}(\pi, \rho, \varphi)$. The lemma is implied by the arbitrariness of ρ and ρ' . $\square$

For a pseudo-metric ρ on $\mathcal{G}^0$, and a sequence $\{\phi_i\}_{i=1}^{\infty}$ with $\phi_i \in [[\mathcal{H}]]_{top}$, a pseudo-metric ρ^{ϕ} on $\mathcal{G}^0$ is given by

$$\rho^\phi(x, y) = \left(\sum_{i=1}^{\infty} 2^{-i} \rho(\phi_i \cdot x, \phi_i \cdot y)^2 \right)^{1/2}.$$

If m is a continuous metric on $\mathcal{G}^0$, then a metric $\overline{m}$ on $[\mathcal{H}]_{top}$ can be given by $\overline{m}(f, g) = \sup_{x \in \mathcal{H}^0} m(f \cdot x, g \cdot x)$. And the topology of $[\mathcal{H}]_{top}$ is independent of the choice of metric m .

Lemma 3.4. [3] *Considering the homeomorphism group of $\mathcal{G}^0$, $Homeo(\mathcal{G}^0)$, with the topology of pointwise convergence, $[\mathcal{H}]_{top}$ is equipped with the topology such that it inherits as a subgroup of $Homeo(\mathcal{G}^0)$. if ρ is a continuous dynamically generating pseudo-metric and $\{\phi_i\}_{i=1}^{\infty}$ is a dense subset of $[\mathcal{H}]_{top}$, then ρ^ϕ is a continuous metric.*

Lemma 3.5. *If ρ is a continuous dynamically generating pseudo-metric and $\{\phi_i\}_{i=1}^{\infty} \subset [\mathcal{H}]_{top}$ with $\phi_1 = \mathcal{H}^0$, then*

$$P_{\mathbb{P}}(\pi, \rho, \varphi) = P_{\mathbb{P}}(\pi, \rho^\phi, \varphi).$$

Proof. For any $x, y \in (\mathcal{G}^0)^d$, it follows from a straightforward computation that $\rho_2^\phi(x, y)^2 = \sum_{j=1}^{\infty} 2^{-j} \rho_2(\phi_j \cdot x, \phi_j \cdot y)^2$. Since $\phi_1 = \mathcal{H}^0$, $\rho \leq 2\rho^\phi$, we have

$$Orb_\nu(\pi, \sigma, F, K, \delta, \rho^\phi) \subset Orb_\nu(\pi, \sigma, F, K, 2\delta, \rho)$$

for any σ, F, K, δ . We claim that

Claim: There exists a $(\rho_2, 2\epsilon)$ -spanning set $\mathcal{E}$ for $Orb_\nu(\pi, \sigma, F, K, 2\delta, \rho)$, which is contained in $Orb_\nu(\pi, \sigma, F, K, 2\delta, \rho)$, and satisfies

$$\begin{aligned} & \sum_{(x_1, \dots, x_d) \in \mathcal{E}} \exp\left(\sum_{i=1}^d \varphi(x_i)\right) \\ & \leq \exp(2\|\varphi\| \lceil \epsilon d \rceil + (d - \lceil \epsilon d \rceil) \eta_1 + 1) \cdot M'_\epsilon(Orb_\nu(\pi, \sigma, F, K, 2\delta, \rho), \rho_2, \varphi), \end{aligned}$$

where $\eta_1 > 0$ such that $|\varphi(x') - \varphi(y')| < \eta_1$, whenever $x', y' \in \mathcal{G}^0$, with $\rho(x', y') < \sqrt{\epsilon}$.

Proof. Let $\mathcal{E}'$ be a minimal (ρ_2, ϵ) -spanning set for $Orb_\nu(\pi, \sigma, F, K, 2\delta, \rho)$, with

$$\sum_{(x'_1, \dots, x'_d) \in \mathcal{E}'} \exp\left(\sum_{i=1}^d \varphi(x'_i)\right) \leq M'_\epsilon(Orb_\nu(\pi, \sigma, F, K, 2\delta, \rho), \rho_2, \varphi) \cdot \exp(1).$$

For each $x' = (x'_1, \dots, x'_d) \in \mathcal{E}'$, there is an element $x = (x_1, \dots, x_d) \in Orb_\nu(\pi, \sigma, F, K, 2\delta, \rho)$ such that $\rho_2(x, x') < \epsilon$. Let $\mathcal{E}$ be the collection of such x , then $\mathcal{E}$ is a $(\rho_2, 2\epsilon)$ -spanning set for $Orb_\nu(\pi, \sigma, F, K, 2\delta, \rho)$. Because $\rho_2(x, x') < \epsilon$, there exists $\Lambda \subset [d]$ such that $|\Lambda| = \lceil \epsilon d \rceil$ and $\rho(x_i, x'_i) < \sqrt{\epsilon}$ for $i \in [d] \setminus \Lambda$. Therefore,

$$\begin{aligned}
& M'_\epsilon(Orb_\nu(\pi, \sigma, F, K, 2\delta, \rho), \rho_2, \varphi) \cdot \exp(1) \\
& \geq \sum_{(x'_1, \dots, x'_d) \in \mathcal{E}'} \exp\left(\sum_{i=1}^d \varphi(x'_i)\right) \\
& \geq \sum_{(x_1, \dots, x_d) \in \mathcal{E}} \exp\left(\sum_{i=1}^d \varphi(x_i) - 2\|\varphi\|\lceil \epsilon d \rceil - (d - \lceil \epsilon d \rceil)\eta_1\right).
\end{aligned}$$

The result is desired. $\square$

Let Ω be the diameter of $(\mathcal{G}^0, \rho)$, F_n be any finite subset of $[[\mathcal{H}]]_{top}$ containing $\{\phi_1, \dots, \phi_n\}$. If $1 \leq i \leq n$ and $x, y \in Orb_\nu(\pi, \sigma, F_n, K, 2\delta, \rho)$ satisfy $\rho_2(x, y) < 2\epsilon$, then

$$\begin{aligned}
& \rho_2(\phi_i \cdot x, \phi_i \cdot y) \\
& \leq \rho_2(x \circ \sigma(\phi_i), y \circ \sigma(\phi_i)) + \rho_2(x \circ \sigma(\phi_i), \phi_i \cdot x) + \rho_2(y \circ \sigma(\phi_i), \phi_i \cdot y) \\
& < 2\epsilon + 4\delta.
\end{aligned}$$

This implies that $\rho_2^\phi(x, y)^2 = \sum_{j=1}^{\infty} 2^{-j} \rho_2(\phi_j \cdot x, \phi_j \cdot y)^2 < 2^{-n} \Omega^2 + (2\epsilon + 4\delta)^2$. By Claim, there exists a $(\rho_2, 2\epsilon)$ -spanning set $\mathcal{E}$ for $Orb_\nu(\pi, \sigma, F_n, K, 2\delta, \rho)$, which is contained in $Orb_\nu(\pi, \sigma, F_n, K, 2\delta, \rho)$, and satisfies

$$\begin{aligned}
& \sum_{(x_1, \dots, x_d) \in \mathcal{E}} \exp\left(\sum_{i=1}^d \varphi(x_i)\right) \\
& \leq \exp(2\|\varphi\|\lceil \epsilon d \rceil + (d - \lceil \epsilon d \rceil)\eta_1 + 1) \cdot M'_\epsilon(Orb_\nu(\pi, \sigma, F_n, K, 2\delta, \rho), \rho_2, \varphi).
\end{aligned}$$

So, for any $x \in Orb_\nu(\pi, \sigma, F_n, K, 2\delta, \rho)$ there exists $y \in \mathcal{E}$ with $\rho_2(x, y) < 2\epsilon$, which implies $\rho_2^\phi(x, y)^2 < 2^{-n} \Omega^2 + (2\epsilon + 4\delta)^2$. Thus, $\mathcal{E}$ is $(\rho_2^\phi, \sqrt{2^{-n} \Omega^2 + (2\epsilon + 4\delta)^2})$ -spanning. Let $\eta = \sqrt{2^{-n} \Omega^2 + (2\epsilon + 4\delta)^2}$. Then

$$\begin{aligned}
& \exp(2\|\varphi\|\lceil \epsilon d \rceil + (d - \lceil \epsilon d \rceil)\eta_1 + 1) \cdot M'_\epsilon(Orb_\nu(\pi, \sigma, F_n, K, 2\delta, \rho), \rho_2, \varphi) \\
& \geq M'_\eta(Orb_\nu(\pi, \sigma, F_n, K, 2\delta, \rho), \rho_2^\phi, \varphi) \\
& \geq M'_\eta(Orb_\nu(\pi, \sigma, F_n, K, \delta, \rho^\phi), \rho_2^\phi, \varphi),
\end{aligned}$$

where the last inequality is due to the inclusion $Orb_\nu(\pi, \sigma, F_n, K, \delta, \rho^\phi) \subset Orb_\nu(\pi, \sigma, F_n, K, 2\delta, \rho)$. By monotonicity, if n is large enough and δ is small enough then $3\epsilon > \eta$ which implies

$$\begin{aligned} & \exp(2\|\varphi\|\lceil \epsilon d \rceil + (d - \lceil \epsilon d \rceil)\eta_1 + 1) \cdot M'_\epsilon(\text{Orb}_\nu(\pi, \sigma, F_n, K, 2\delta, \rho), \rho_2, \varphi) \\ & \geq M'_{3\epsilon}(\text{Orb}_\nu(\pi, \sigma, F_n, K, \delta, \rho^\phi), \rho_2^\phi, \varphi). \end{aligned}$$

Since F_n is any finite subset of $[[\mathcal{H}]]_{\text{top}}$ containing $\{\phi_1, \dots, \phi_n\}$ and Remark 3.2, we have

$$\begin{aligned} & 2\|\varphi\|\epsilon + (1 - \epsilon)\eta_1 + \inf_{\delta > 0} \inf_{F \subset_f [[\mathcal{H}]]_{\text{top}}} \inf_{K \subset_f C(\mathcal{H}^0)} \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|M'_\epsilon(\text{Orb}_\nu(\pi, \cdot, F, K, 2\delta, \rho), \rho_2, \varphi)\|_{p, \mathbb{P}_j} \\ & \geq \inf_{\delta > 0} \inf_{F \subset_f [[\mathcal{H}]]_{\text{top}}} \inf_{K \subset_f C(\mathcal{H}^0)} \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|M'_{3\epsilon}(\text{Orb}_\nu(\pi, \cdot, F, K, \delta, \rho^\phi), \rho_2^\phi, \varphi)\|_{p, \mathbb{P}_j}. \end{aligned}$$

Since this is true for every $\epsilon > 0$ and $\eta_1 \rightarrow 0$ as $\epsilon \rightarrow 0$, we have $P_\mathbb{P}(\pi, \rho, 2, \varphi) \geq P_\mathbb{P}(\pi, \rho^\phi, 2, \varphi)$, which implies $P_\mathbb{P}(\pi, \rho, \varphi) \geq P_\mathbb{P}(\pi, \rho^\phi, \varphi)$.

Now, it's turn to show the opposite inequality. It was claimed that given any finite $F \subset [[\mathcal{H}]]_{\text{top}}$ with $\mathcal{H}^0 \in F$ and $\delta > 0$, if n is sufficiently large, $F' = \{\phi_j f : f \in F, 1 \leq j \leq n\}$ and σ is $(F', \delta^2/M^2)$ -multiplicative then

$$\text{Orb}_\nu(\pi, \sigma, F, K, 2\delta, \rho^\phi) \supset \text{Orb}_\nu(\pi, \sigma, F', K, \delta, \rho)$$

in [3]. The claim and $\rho \leq 2\rho^\phi$ show that

$$M'_\epsilon(\text{Orb}_\nu(\pi, \sigma, F, K, 2\delta, \rho^\phi), \rho_\infty^\phi, \varphi) \geq M'_{2\epsilon}(\text{Orb}_\nu(\pi, \sigma, F', K, \delta, \rho), \rho_\infty, \varphi).$$

This implies that $P_\mathbb{P}(\pi, \rho^\phi, \varphi) \geq P_\mathbb{P}(\pi, \rho, \varphi)$. □

Proof of Theorem 3.1 By [3], there exists a dense sequence $\phi = \{\phi_i\}_{i=1}^\infty \subset [\mathcal{H}]_{\text{top}}$ with $\phi_1 = \mathcal{H}^0$. By Lemmas 3.5, 3.4, 3.3, $P_\mathbb{P}(\pi, \rho, \varphi) = P_\mathbb{P}(\pi, \rho^\phi, \varphi) = P_\mathbb{P}(\pi, \rho'^\phi, \varphi) = P_\mathbb{P}(\pi, \rho', \varphi)$. □

4 Measure entropy

There are two definitions of measure sofic entropy for groupoid. One is defined via partitions by Bowen [3] in a manner analogous to [12], the other is via pseudo-metric like topological entropy.

Let $\pi : (\mathcal{G}, \mu) \rightarrow (\mathcal{H}, \nu)$ be a class-bijective extension of pmp discrete groupoids and $\mathbb{P} := \{\mathbb{P}_j\}_{j \in J}$ a sofic approximation to $(\mathcal{H}, \nu)$. For a finite partition $\mathcal{P}$ of $\mathcal{G}^0$ and $F \subset_f [[\mathcal{H}]]$, let $\mathcal{P}^F$ be the coarsest partition of $\mathcal{G}^0$ containing $\{f \cdot P : f \in F, P \in \mathcal{P}\}$. Let $\Sigma(\mathcal{P})$ denote the smallest sigma algebra of $\mathcal{G}^0$ containing $\mathcal{P}$. Let $\mathcal{B}(\Delta_d^0)$ be the collection of all subsets of Δ_d^0 . A map $\phi : \Sigma(\mathcal{P}) \rightarrow \mathcal{B}(\Delta_d^0)$ is a homeomorphism means that given $A, B \in \Sigma(\mathcal{P})$, $\phi(A \cup B) = \phi(A) \cup \phi(B)$, $\phi(A \cap B) = \phi(A) \cap \phi(B)$ and $\phi(\mathcal{G}^0) = \Delta_d^0$.

Definition 4.1. [3](Good homomorphism) For $\sigma : [[\mathcal{H}]] \rightarrow [[d]], f \in [[\mathcal{H}]], \delta > 0$ and $F \subset [[\mathcal{H}]]$, let $Hom(\pi, \sigma, \mathcal{P}, F, \delta)$ be the collection of all homomorphisms $\phi : \Sigma(\mathcal{P}^F) \rightarrow \mathcal{B}(\Delta_d^0)$ such that

- (1) $\sum_{P \in \mathcal{P}} |\sigma(f) \cdot \phi(P) \Delta \phi(f \cdot P)| < d\delta, \forall f \in F;$
- (2) $\sum_{P \in \mathcal{P}^F} |\phi(P)|d^{-1} - \mu(P)| < \delta.$

Given a partition $\mathcal{Q}$ of $\mathcal{G}^0$ with $\mathcal{Q} \leq \mathcal{P}$, let $|Hom(\pi, \sigma, \mathcal{P}, F, \delta)|_{\mathcal{Q}}$ be the cardinality of the set of homomorphisms $\phi : \Sigma(\mathcal{Q}) \rightarrow \mathcal{B}(\Delta_d^0)$ such that there exists a $\phi' \in Hom(\pi, \sigma, \mathcal{P}, F, \delta)$ satisfying that ϕ is the restriction of ϕ' to $\Sigma(\mathcal{Q})$.

Definition 4.2. [3] Let $\mathcal{B}(\mathcal{G}^0)$ be the Borel sigma algebra of $\mathcal{G}^0$. For finite Borel partitions $\mathcal{Q} \leq \mathcal{P}$ and a sub-algebra $\mathcal{F} \subset \mathcal{B}(\mathcal{G}^0)$, define

$$\begin{aligned} h_{\mathbb{P}, \mu}(\pi, \mathcal{Q}, \mathcal{P}, F, \delta) &:= \lim_{j \rightarrow \beta} \frac{1}{d_j} \log ||Hom(\pi, \cdot, \mathcal{P}, F, \delta)|_{\mathcal{Q}}||_{p, \mathbb{P}_j} \\ h_{\mathbb{P}, \mu}(\pi, \mathcal{Q}, \mathcal{P}) &:= \inf_{F \subset_f [[\mathcal{H}]]} \inf_{\delta > 0} h_{\mathbb{P}, \mu}(\pi, \mathcal{Q}, \mathcal{P}, F, \delta) \\ h_{\mathbb{P}, \mu}(\pi, \mathcal{Q}, \mathcal{F}) &:= \inf_{\mathcal{Q} \leq \mathcal{P} \subset \mathcal{F}} h_{\mathbb{P}, \mu}(\pi, \mathcal{Q}, \mathcal{P}) \\ h_{\mathbb{P}, \mu}(\pi, \mathcal{F}) &:= \sup_{\mathcal{Q} \subset \mathcal{F}} h_{\mathbb{P}, \mu}(\pi, \mathcal{Q}, \mathcal{F}). \end{aligned}$$

The infimum in the second-to-last line runs over all finite Borel partitions $\mathcal{P}$ with $\mathcal{Q} \leq \mathcal{P} \subset \mathcal{F}$ and the supremum in the last line runs over all finite partition $\mathcal{Q} \subset \mathcal{F}$. The sofic measure entropy of π with respect to $\mathbb{P}, p, \beta$ is given by $h_{\mathbb{P}, \mu}(\pi) := h_{\mathbb{P}, \mu}(\pi, \mathcal{B}(\mathcal{G}^0))$.

Remark 4.1. [3]

- $h_{\mathbb{P}, \mu}(\pi)$ depends implicitly on $1 \leq p \leq \infty$ and a bias β .
- The order of the supremums, infimums and limits above is important with the exception that the three infimums can be permuted without affecting the definition of $h_{\mathbb{P}, \mu}(\pi, \mathcal{Q}, \mathcal{F})$.
- The quantity $\frac{1}{d_j} \log ||Hom(\pi, \cdot, \mathcal{P}, F, \delta)|_{\mathcal{Q}}||_{p, \mathbb{P}_j}$

is monotone increasing in $\delta, \mathcal{Q}$ and decreasing in $F, \mathcal{P}$.

Definition 4.3. [3] Let $(\mathcal{H}, \nu)$ be a discrete pmp groupoid, $\sigma : [[\mathcal{H}]] \rightarrow [[d]], F \subset_f [[\mathcal{H}]]$ and $\delta > 0$. σ is (F, δ) -continuous means that $|\sigma(f) \Delta \sigma(g)|_d < \delta + \nu(f \Delta g), \forall f, g \in F$. If $\mathbb{P} = \{\mathbb{P}_j\}_{j \in J}$ is a sofic approximation to $(\mathcal{H}, \nu)$, then $\mathbb{P}$ is asymptotically continuous means that for every $F \subset_f [[\mathcal{H}]], \delta > 0$, there exists $j \in J$ such that $j' \geq j$ implies $\mathbb{P}_{j'}$ -almost every σ is (F, δ) -continuous.

Now, let's present the formulation of measure entropy via pseudo-metrics.

The assumption is given as follows: two discrete pmp topological groupoids $\mathcal{G}, \mathcal{H}$ such that $\mathcal{G}^0$ and $\mathcal{H}^0$ are compact metrizable spaces, a class-bijective continuous factor $\pi : (\mathcal{G}, \mu) \rightarrow (\mathcal{H}, \nu)$, a sofic approximation $\mathbb{P} = \{\mathbb{P}_j\}_{j \in J}$ to $(\mathcal{H}, \nu)$, a continuous pseudo-metric ρ on $\mathcal{G}^0$, a bias β and $p \in [1, \infty]$. From this data, the sofic measure entropy of (π, ρ) with respect to $(\mathbb{P}, p, \beta)$ can be defined. Furthermore, if ρ is dynamically generating, $\mathcal{H}$ is étale, ν is regular and $\mathbb{P}$ is asymptotically continuous, then this entropy coincides with the Definition 4.2. The reader is referred to [3] for details.

Definition 4.4. [3] *For a map $\sigma : [[\mathcal{H}]] \rightarrow [[d]], F \subset_f [[\mathcal{H}]]_{top}, K \subset_f C(\mathcal{G}^0)$ and $\delta > 0$, let $Orb_\mu(\pi, \sigma, F, K, \delta, \rho)$ be the set of all $(x_1, \dots, x_d) \in Orb_\nu(\pi, \sigma, F, \emptyset, \delta, \rho)$ such that*

$$\max_{x \in K} \left| \frac{1}{d} \sum_{i=1}^d k(x_i) - \int_{\mathcal{G}^0} k d\mu \right| < \delta.$$

Define

$$h_{\mathbb{P}, \mu}(\pi, \rho, 2) := \sup_{\epsilon > 0} \inf_{\delta > 0} \inf_{F \subset_f [[\mathcal{H}]]_{top}} \inf_{K \subset_f C(\mathcal{G}^0)} \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|N_\epsilon(Orb_\mu(\pi, \cdot, F, K, \delta, \rho), \rho_2)\|_{p, \mathbb{P}_j},$$

$$h_{\mathbb{P}, \mu}(\pi, \rho, \infty) := \sup_{\epsilon > 0} \inf_{\delta > 0} \inf_{F \subset_f [[\mathcal{H}]]_{top}} \inf_{K \subset_f C(\mathcal{G}^0)} \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|N_\epsilon(Orb_\mu(\pi, \cdot, F, K, \delta, \rho), \rho_\infty)\|_{p, \mathbb{P}_j}.$$

Remark 4.2. [3]

- $h_{\mathbb{P}, \mu}(\pi, \rho, 2)$ and $h_{\mathbb{P}, \mu}(\pi, \rho, \infty)$ depend implicitly on the choice of bias β and parameter $p \in [1, \infty]$.
- The order of the supremums, infimums and limits above is important with the exception that the three infimums can be permuted without affecting the definition.
- The quantity $\frac{1}{d_j} \log \|N_\epsilon(Orb_\mu(\pi, \cdot, F, K, \delta, \rho), \rho_2)\|_{p, \mathbb{P}_j}$ is monotone increasing in δ and decreasing in ϵ, F, K .
- Similar statements remain valid if ρ_2 is replaced with ρ_∞ or N_ϵ is replaced with N'_ϵ .
- If we replace $N_\epsilon(\cdot)$ with $N'_\epsilon(\cdot)$, then we get equivalent definitions.
- $h_{\mathbb{P}, \mu}(\pi, \rho, 2) = h_{\mathbb{P}, \mu}(\pi, \rho, \infty)$.
- Let $h_{\mathbb{P}, \mu}(\pi, \rho) := h_{\mathbb{P}, \mu}(\pi, \rho, 2)$. If ρ, ρ' are dynamically generating continuous pseudo-metrics on $\mathcal{G}^0$ then $h_{\mathbb{P}, \mu}(\pi, \rho) = h_{\mathbb{P}, \mu}(\pi, \rho')$.
- If ρ is a dynamically generating continuous pseudo-metric on $\mathcal{G}^0$, $\mathcal{H}$ is étale, ν is regular and $\mathbb{P}$ is asymptotically continuous then $h_{\mathbb{P}, \mu}(\pi, \rho) = h_{\mathbb{P}, \mu}(\pi)$.

5 The variational principle

This section is aim to reveal the relationship between topological pressure and measure entropy. First, we present two propositions the proof of which can be found in [3].

Proposition 5.1. [3] *Let $(\mathcal{H}, \nu)$ be a pmp separable étale topological discrete groupoid, $\mathcal{G}$ be a separable topological discrete groupoid, and μ be a Borel probability measure on $\mathcal{G}^0$, and suppose that μ is $[[\mathcal{H}]]_{top}$ -invariant in the sense that $\mu(k \circ f) = \mu(k \circ \tau(f))$ for any continuous function $k \in C(\mathcal{G}^0)$ and $f \in [[\mathcal{H}]]_{top}$, then $(\mathcal{G}, \mu)$ is pmp.*

Proposition 5.2. [3] *Assume that Ω is a directed set and $\omega \in \Omega \mapsto \mu_\omega \in M(\mathcal{G}^0)$ is a map such that $\lim_{\omega \rightarrow \Omega} \mu_\omega = \mu_\infty$, and $\pi_* \mu_\infty = \nu$. Then $\lim_{\omega \rightarrow \Omega} \mu_\omega(k \circ f) = \mu_\infty(k \circ f)$, for any $k \in C(\mathcal{G}^0)$ and $f \in [[\mathcal{H}]]_{top}$.*

Theorem 5.1. *Let $(\mathcal{H}, \nu)$ be a pmp separable étale topological discrete groupoid, $\mathcal{G}$ be a separable topological discrete groupoid, $\pi : \mathcal{G} \rightarrow \mathcal{H}$ a continuous class-bijective factor, and $\mathbb{P} = \{\mathbb{P}_j\}_{j \in J}$ an asymptotically continuous sofic approximation to $(\mathcal{H}, \nu)$. If $\mathcal{H}^0$ and $\mathcal{G}^0$ are compact and metrizable and ν is regular, then for any $\varphi \in C(\mathcal{G}^0, \mathbb{R})$, $p \in [1, \infty]$ and bias $\beta \neq -$,*

$$P_{\mathbb{P}}(\pi, \varphi) = \max_{\mu} h_{\mathbb{P}, \mu}(\pi) + \int_{\mathcal{G}^0} \varphi d\mu,$$

where the max runs over all measures μ on $\mathcal{G}^0$ such that $\pi_* \mu = \nu$ and $(\mathcal{G}, \mu)$ is pmp.

Proof. The proof will be divided into the following two steps.

Step 1: This step proves $P_{\mathbb{P}}(\pi, \varphi) \leq \sup_{\mu} h_{\mathbb{P}, \mu}(\pi) + \int_{\mathcal{G}^0} \varphi d\mu$.

Without loss of generality, we may assume $P_{\mathbb{P}}(\pi, \varphi) > -\infty$. Let $\kappa > 0$, then there exists $\epsilon > 0$ such that

$$P_{\mathbb{P}}^{\epsilon}(\pi, \rho, 2, \varphi) \geq P_{\mathbb{P}}(\pi, \rho, 2, \varphi) - \kappa,$$

where $P_{\mathbb{P}}^{\epsilon}(\pi, \rho, 2, \varphi) := \inf_{\delta > 0} \inf_{F \subset_f [[\mathcal{H}]]_{top}} \inf_{K \subset_f C(\mathcal{H}^0)} \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|M_{\epsilon}(Orb_{\nu}(\pi, \cdot, F, K, \delta, \rho), \rho_2, \varphi)\|_{p, \mathbb{P}_j}$.

Let $\Omega = \{(F, L, \delta) : F \subset_f [[\mathcal{H}]]_{top}, L \subset_f C(\mathcal{G}^0), \delta > 0\}$. Ω is considered as a directed set by declaring $(F, L, \delta) \leq (F', L', \delta')$ if $F' \supset F, L' \supset L, \delta' \leq \delta$. Given $\omega = (F_{\omega}, L_{\omega}, \delta_{\omega}) \in \Omega$, let $K_{\omega} = \{k \in C(\mathcal{H}^0) : k \circ \pi \in L_{\omega}\}$. Let $M(\mathcal{G}^0)$ denote the space of Borel probability measures on $\mathcal{G}^0$.

Claim: There exists a directed net $\omega \in \Omega \mapsto \mu_{\omega} \in M(\mathcal{G}^0)$ such that

(i) $h_{\mathbb{P}, \mu_{\omega}}^{\epsilon}(\pi, \rho, \omega, 2) + \int_{\mathcal{G}^0} \varphi d\mu_{\omega} + \delta_{\omega} \geq P_{\mathbb{P}}^{\epsilon}(\pi, \rho, 2, \varphi)$, where

$$\begin{aligned} h_{\mathbb{P}, \mu_{\omega}}^{\epsilon}(\pi, \rho, \omega, 2) &:= h_{\mathbb{P}, \mu_{\omega}}^{\epsilon}(\pi, \rho, F_{\omega}, L_{\omega}, \delta_{\omega}, 2) \\ &:= \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|N_{\epsilon}(Orb_{\mu_{\omega}}(\pi, \cdot, F_{\omega}, L_{\omega}, \delta_{\omega}, \rho), \rho_2)\|_{p, \mathbb{P}_j}. \end{aligned}$$

(ii) $\lim_{\omega \rightarrow \Omega} |\mu_\omega(k \circ f) - \mu_\omega(k \circ \tau(f))| = 0$, for any $f \in [[\mathcal{H}]]_{top}$, $k \in C(\mathcal{G}^0)$.

(iii) $\lim_{\omega \rightarrow \Omega} |\mu_\omega(k \circ \pi) - \nu(k)| = 0$, for any $k \in C(\mathcal{H}^0)$.

Proof. For $\varphi \in L \subset_f C(\mathcal{G}^0)$ and $\delta > 0$, let $M(L, \delta)$ be the set of all $\mu \in M(\mathcal{G}^0)$ such that $|\mu(k \circ \pi) - \nu(k)| \leq \delta$ for any $k \in C(\mathcal{H}^0)$ with $k \circ \pi \in L$. For $F \subset_f [[\mathcal{H}]]_{top}$, let $D(F, L, \delta) \subset M(L, \delta)$ be a finite set satisfying that for every $\lambda \in M(L, \delta)$ there exists a $\mu \in D(F, L, \delta)$ such that

$$|\mu(k \circ f) - \lambda(k \circ f)| < \delta, \quad \forall f \in F, k \in L.$$

For every $x \in (\mathcal{G}^0)^d$, the measure $m_x \in M(\mathcal{G}^0)$ is given by $m_x = d^{-1} \sum_{i=1}^d \delta_{x_i}$, where δ_{x_i} is the Dirac measure concentrated on x_i . For every $\omega \in \Omega$, choose a Borel map $x \in \{y \in (\mathcal{G}^0)^d : m_y \in M(L_\omega, \delta_\omega)\} \mapsto \mu_{x,\omega} \in D(\omega)$ such that

$$|\mu_{x,\omega}(k \circ f) - m_x(k \circ f)| < \delta_\omega, \quad \forall f \in F_\omega, k \in L_\omega.$$

For any $\sigma : [[\mathcal{H}]] \rightarrow [[d]]$, $F \subset_f [[\mathcal{H}]]_{top}$, $K \subset_f C(\mathcal{H}^0)$ and $\delta > 0$, choose a maximum (ρ_2, ϵ) -separated subset $Q(\sigma, F, K, \delta) \subset Orb_\nu(\pi, \sigma, F, K, \delta, \rho)$ such that for any d_j , the map $\sigma \in \text{Map}([[\mathcal{H}]], [[d_j]]) \mapsto Q(\sigma, F, K, \delta)$ is Borel, and

$$\exp(1) \sum_{(x_1, \dots, x_d) \in Q(\sigma, F, K, \delta)} \exp\left(\sum_{i=1}^d \varphi(x_i)\right) \geq M_\epsilon(Orb_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_2, \varphi).$$

By the Pigeonhole Principle for any $\omega \in \Omega$ and $j \in J$ there exists $\mu_{j,\omega} \in D(\omega)$ such that

$$\begin{aligned} & \left\| \sum_{(x_1, \dots, x_d) \in \{x \in Q(\cdot, F_\omega, K_\omega, \delta_\omega) : \mu_{j,\omega} = \mu_{x,\omega}\}} \exp\left(\sum_{i=1}^d \varphi(x_i)\right) \right\|_{p, \mathbb{P}_j} \\ & \geq \frac{\left\| \sum_{(x_1, \dots, x_d) \in Q(\cdot, F_\omega, K_\omega, \delta_\omega)} \exp\left(\sum_{i=1}^d \varphi(x_i)\right) \right\|_{p, \mathbb{P}_j}}{|D(\omega)|} \\ & \geq \frac{\|M_\epsilon(Orb_\nu(\pi, \cdot, F_\omega, K_\omega, \delta_\omega, \rho), \rho_2, \varphi)\|_{p, \mathbb{P}_j}}{|D(\omega)|}. \end{aligned}$$

Then choose $\mu_\omega \in D(\omega)$ so that if $J' = \{j \in J : \mu_{j,\omega} = \mu_\omega\}$ then either $J' \in \beta$ (if β is an ultrafilter on J) or J' is cofinal. If $\beta = +$, we also require that

$$\begin{aligned} & \limsup_{j \in J} d_j^{-1} \log \|M_\epsilon(Orb_{\mu_{j,\omega}}(\pi, \cdot, F_\omega, L_\omega, \delta_\omega), \rho_2, \varphi)\|_{p, \mathbb{P}_j} \\ & = \limsup_{j \in J'} d_j^{-1} \log \|M_\epsilon(Orb_{\mu_{j,\omega}}(\pi, \cdot, F_\omega, L_\omega, \delta_\omega), \rho_2, \varphi)\|_{p, \mathbb{P}_j}. \end{aligned}$$

This is possible due to the finiteness of $D(\omega)$. The measures $\{\mu_\omega : \omega \in \Omega\}$ is desired. First, for any $j \in J, \omega \in \Omega$, if $x \in Orb_\nu(\pi, \sigma, F_\omega, K_\omega, \delta_\omega, \rho)$ and $\mu_{x,\omega} = \mu_{j,\omega}$, then

$$|m_x(k) - \mu_{j,\omega}(k)| < \delta_\omega, \quad \forall k \in L_\omega$$

implies $x \in Orb_{\mu_{j,\omega}}(\pi, \sigma, F_\omega, L_\omega, \delta_\omega, \rho)$. Therefore,

$$\begin{aligned} & \sum_{(x_1, \dots, x_d) \in \{x \in Q(\sigma, F_\omega, K_\omega, \delta_\omega) : \mu_{j,\omega} = \mu_{x,\omega}\}} \exp\left(\sum_{i=1}^d \varphi(x_i)\right) \\ & \leq \sum_{(x_1, \dots, x_d) \in \{x \in Q(\sigma, F_\omega, K_\omega, \delta_\omega) : \mu_{j,\omega} = \mu_{x,\omega}\}} \exp(d\mu_{j,\omega}(\varphi) + d\delta_\omega) \\ & = |\{x \in Q(\sigma, F_\omega, K_\omega, \delta_\omega) : \mu_{j,\omega} = \mu_{x,\omega}\}| \cdot \exp(d\mu_{j,\omega}(\varphi) + d\delta_\omega) \\ & \leq N_\epsilon(Orb_{\mu_{j,\omega}}(\pi, \sigma, F_\omega, L_\omega, \delta_\omega, \rho), \rho_2) \cdot \exp(d\mu_{j,\omega}(\varphi) + d\delta_\omega). \end{aligned}$$

By choice of $\mu_{j,\omega}$, this implies

$$\begin{aligned} & \|N_\epsilon(Orb_{\mu_{j,\omega}}(\pi, \cdot, F_\omega, L_\omega, \delta_\omega, \rho), \rho_2)\|_{p, \mathbb{P}_j} \cdot \exp(d\mu_{j,\omega}(\varphi) + d\delta_\omega) \\ & \geq \frac{\|M_\epsilon(Orb_\nu(\pi, \cdot, F_\omega, K_\omega, \delta_\omega, \rho), \rho_2, \varphi)\|_{p, \mathbb{P}_j}}{|D(\omega)|} \exp(-1). \end{aligned}$$

Due to the choice of μ_ω , it follows that

$$\begin{aligned} & h_{\mathbb{P}, \mu_\omega}^\epsilon(\pi, \rho, \omega, 2) \\ & \geq \lim_{j \rightarrow \beta} d_j^{-1} \log \|N_\epsilon(Orb_{\mu_{j,\omega}}(\pi, \cdot, F_\omega, L_\omega, \delta_\omega, \rho), \rho_2)\|_{p, \mathbb{P}_j} \\ & \geq \lim_{j \rightarrow \beta} d_j^{-1} \log \|M_\epsilon(Orb_\nu(\pi, \cdot, F_\omega, L_\omega, \delta_\omega, \rho), \rho_2, \varphi)\|_{p, \mathbb{P}_j} - \mu_\omega(\varphi) - \delta_\omega \\ & \geq P_{\mathbb{P}}^\epsilon(\pi, \rho, 2, \varphi) - \mu_\omega(\varphi) - \delta_\omega. \end{aligned}$$

$\beta \neq -$ is used in the first inequality. This finishes the proof of the first item of Claim.

The proof of the following two items can be seen in [3]. We provide it here for completeness. To prove the second item, let $k \in C(\mathcal{G}^0)$, $f \in [[\mathcal{H}]]_{top}$ and $\eta > 0$ be a constant. Since k is continuous, there exists a constant $\delta > 0$ such that if $x, y \in \mathcal{G}^0$ with $\rho(x, y) < \delta$, then $|k(x) - k(y)| < \eta$.

Let $\omega \in \Omega$ be such that $k \in L_\omega$, $f, f^{-1}, \tau(f) \in F_\omega$ and δ_ω is small enough so that $\delta_\omega^2/\delta^2 < \eta$. Due to the choice of μ_ω , there exist a $(F_\omega, \delta_\omega^2/(100\text{diam}(\rho)^2))$ -multiplicative $\sigma : [[\mathcal{H}]] \rightarrow [[d]]$ and $x \in Orb_\nu(\pi, \sigma, F_\omega, K_\omega, \delta_\omega, \rho)$ such that $\mu_{x,\omega} = \mu_\omega$. It follows from a straightforward computation that $|\mu_\omega(k \circ f) - \mu_\omega(k \circ \tau(f))| \leq 2\delta_\omega + |m_x(k \circ f) - m_{x \circ \sigma(f)}(k)| + |m_{x \circ \sigma(f)}(k) - m_x(k \circ \tau(f))|$. It follows from $x \in Orb_\nu(\pi, \sigma, F_\omega, K_\omega, \delta_\omega, \rho)$ that

$$\begin{aligned}\delta_\omega^2 &\geq \rho_2(f \cdot x, x \circ \sigma(f))^2 \\ &\geq d^{-1} |\{1 \leq i \leq d : \rho(f \cdot x_i, x_{\sigma(f)i}) \geq \delta\}| \delta^2.\end{aligned}$$

So, $\eta > \frac{\delta_\omega^2}{\delta^2} \geq d^{-1} |\{1 \leq i \leq d : \rho(f \cdot x_i, x_{\sigma(f)i}) \geq \delta\}|$ which implies

$$|m_x(k \circ f) - m_{x \circ \sigma(f)}(k)| \leq d^{-1} \sum_{i=1}^d |k(f \cdot x_i) - k(x_{\sigma(f)i})| \leq 2\eta + 2\eta \|k\|.$$

It follows from $x \in Orb_\nu(\pi, \sigma, F_\omega, K_\omega, \delta_\omega, \rho)$ that

$$\begin{aligned}\delta_\omega^2 &> \rho_2(x \circ \sigma(\tau(f)), \tau(f) \cdot x)^2 = d^{-1} \sum_{i=1}^d \rho(x_{\sigma(\tau(f))i}, \tau(f) \cdot x_i)^2 \\ &\geq d^{-1} \text{diam}(\rho)^2 |\tau(\sigma(\tau(f))) \Delta \{i : x_i \in \tau(f)\}| \\ &\geq d^{-1} \text{diam}(\rho)^2 |\tau(\sigma(f)) \Delta \{i : x_i \in \tau(f)\}| - \delta_\omega^2.\end{aligned}$$

This implies that

$$\begin{aligned}&|m_{x \circ \sigma(f)}(k) - m_x(k \circ \tau(f))| \\ &= d^{-1} \left| \sum_{i \in \varsigma(\sigma(f))} k(x_{\sigma(f)i}) - \sum_{i: x_i \in \tau(f)} k(x_i) \right| \\ &= d^{-1} \left| \sum_{i \in \tau(\sigma(f))} k(x_{\sigma(f)i}) - \sum_{i: x_i \in \tau(f)} k(x_i) \right| \\ &\leq d^{-1} \|k\| \cdot |\tau(\sigma(f)) \Delta \{i : x_i \in \tau(f)\}| \\ &\leq d^{-1} \|k\| \frac{2\delta_\omega^2}{d^{-1} \text{diam}(\rho)^2} = \frac{2\delta_\omega^2 \|k\|}{\text{diam}(\rho)^2}.\end{aligned}$$

Hence,

$$|\mu_\omega(k \circ f) - \mu_\omega(k \circ \tau(f))| \leq 2\delta_\omega + 2\eta + 2\eta \|k\| + \frac{2\delta_\omega^2 \|k\|}{\text{diam}(\rho)^2}.$$

Due to the arbitrariness of η, f, k , we obtain

$$\lim_{\omega \rightarrow \Omega} |\mu_\omega(k \circ f) - \mu_\omega(k \circ \tau(f))| = 0,$$

for any $f \in [[\mathcal{H}]]_{top}, k \in C(\mathcal{G}^0)$.

It's turn to show the proof of the third of Claim. Let $k \in C(\mathcal{H}^0)$. let $\omega \in \Omega$ be such that $k \in K_\omega$. Due to the choice of μ_ω , there exists $\sigma : [[\mathcal{H}]] \rightarrow [[d]]$ and $x \in Orb_\nu(\pi, \sigma, F_\omega, K_\omega, \delta_\omega, \rho)$ with $\mu_{x,\omega} = \mu_\omega$. This implies that

$$|\mu_\omega(k \circ \pi) - \nu(k)| \leq |\mu_\omega(k \circ \pi) - m_x(k \circ \pi)| + |m_x(k \circ \pi) - \nu(k)| < 2\delta_\omega.$$

Therefore, $\lim_{\omega \rightarrow \Omega} |\mu_\omega(k \circ \pi) - \nu(k)| = 0$ as required. The proof of Claim is completed. $\square$

Let μ be a weak* accumulation point of $\{\mu_\omega : \omega \in \Omega\}$. It follows from the third item of Claim that $\pi_*\mu = \nu$. Due to (ii) and Proposition 5.2, we have $\mu(k \circ f) = \mu(k \circ \tau(f))$ for any $f \in [[\mathcal{H}]]_{top}$, $k \in \mathcal{G}^0$. By Proposition 5.1, $(\mathcal{G}, \mu)$ is pmp. Let $F \subset_f [[\mathcal{H}]]_{top}$, $\varphi \in L \subset_f C(\mathcal{G}^0)$ and $\delta > 0$. Choose $\omega \in \Omega$ to satisfy

- (1) $|\mu(k) - \mu_\omega(k)| \leq \delta/2, \forall k \in L$;
- (2) $F \subset F_\omega, L \subset L_\omega, \delta_\omega \leq \delta/2$.

Then for any $\sigma : [[\mathcal{H}]] \rightarrow [[d]]$, $x \in Orb_{\mu_\omega}(\pi, \sigma, F_\omega, L_\omega, \delta_\omega, \rho)$ and $k \in L$,

$$\left| \frac{1}{d} \sum_{i=1}^d k(x_i) - \mu(k) \right| \leq \left| \frac{1}{d} \sum_{i=1}^d k(x_i) - \mu_\omega(k) \right| + |\mu_\omega(k) - \mu(k)| < \delta,$$

which implies that

$$Orb_{\mu_\omega}(\pi, \sigma, F_\omega, L_\omega, \delta_\omega, \rho) \subset Orb_\mu(\pi, \sigma, F, L, \delta, \rho).$$

So,

$$\begin{aligned} & h_{\mathbb{P},\mu}^\epsilon(\pi, \rho, F, L, \delta, 2) + \mu(\varphi) + \delta \\ & \geq h_{\mathbb{P},\mu_\omega}^\epsilon(\pi, \rho, F_\omega, L_\omega, \delta_\omega, 2) + \mu_\omega(\varphi) \\ & \geq P_{\mathbb{P}}^\epsilon(\pi, \rho, 2, \varphi) - \delta_\omega \\ & \geq P_{\mathbb{P}}^\epsilon(\pi, \rho, 2, \varphi) - \delta/2. \end{aligned}$$

By taking the infimum over F, L, δ , we get

$$\begin{aligned} & h_{\mathbb{P},\mu}(\pi, \rho, 2) + \mu(\varphi) \geq h_{\mathbb{P},\mu}^\epsilon(\pi, \rho, 2) + \mu(\varphi) \\ & \geq P_{\mathbb{P}}^\epsilon(\pi, \rho, 2, \varphi) \geq P_{\mathbb{P}}(\pi, \rho, 2, \varphi) - \kappa. \end{aligned}$$

Step 1 follows due to the arbitrariness of κ .

Step 2: This step proves $P_{\mathbb{P}}(\pi, \varphi) \geq \max_{\mu} h_{\mathbb{P},\mu}(\pi) + \int_{\mathcal{G}^0} \varphi d\mu$.

Fix $\mu \in M(\mathcal{G}^0)$ with $\pi_*\mu = \nu$. Let $\delta > 0$, $F \subset_f [[\mathcal{H}]]_{top}$ and $K \subset_f C(\mathcal{H}^0)$. Let $\{\varphi\} \cup \{k \circ \pi : k \in K\} \subset K' \subset_f C(\mathcal{G}^0)$. Then for any $\sigma : [[\mathcal{H}]] \rightarrow [[d]]$,

$$Orb_\mu(\pi, \sigma, F, K', \delta, \rho) \subset Orb_\nu(\pi, \sigma, F, K, \delta, \rho).$$

Then

$$\begin{aligned} & M_\epsilon(Orb_\nu(\pi, \sigma, F, K, \delta, \rho), \rho_\infty, \varphi) \\ & \geq N_\epsilon(Orb_\mu(\pi, \sigma, F, K', \delta, \rho), \rho_\infty) \cdot \exp(d\mu(\varphi) - d\delta - 1). \end{aligned}$$

This implies that

$$\begin{aligned} & \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|M_\epsilon(Orb_\nu(\pi, \cdot, F, K, \delta, \rho), \rho_\infty, \varphi)\|_{p, \mathbb{P}_j} \\ & \geq \lim_{j \rightarrow \beta} \frac{1}{d_j} \log \|N_\epsilon(Orb_\mu(\pi, \sigma, F, K', \delta, \rho), \rho_\infty)\|_{p, \mathbb{P}_j} + \mu(\varphi) - \delta. \end{aligned}$$

By taking the infimum over F, K', K, δ and the supremum over ϵ , then $P_{\mathbb{P}}(\pi, \varphi) \geq h_{\mathbb{P}, \mu}(\pi) + \int_{\mathcal{G}^0} \varphi d\mu$. This finishes the proof. $\square$

Some properties of topological pressure are presented as below without proof. It is easier to investigate these properties of topological pressure by variational principle than by definition.

Proposition 5.3. *The conditions are given as the conditions of Theorem 5.1.*

- (i) $P_{\mathbb{P}}(\pi, \mathbf{0})$ is topological entropy defined in [3].
- (ii) $P_{\mathbb{P}}(\pi, \varphi + c) = P_{\mathbb{P}}(\pi, \varphi) + c$, where c is a constant.
- (iii) $P_{\mathbb{P}}(\pi, \varphi + \psi) \leq P_{\mathbb{P}}(\pi, \varphi) + P_{\mathbb{P}}(\pi, \psi)$, where $\psi \in C(\mathcal{G}^0, \mathbb{R})$.
- (iv) $\varphi \leq \psi$ implies $P_{\mathbb{P}}(\pi, \varphi) \leq P_{\mathbb{P}}(\pi, \psi)$. In particular, $P_{\mathbb{P}}(\pi, \mathbf{0}) + \min \varphi \leq P_{\mathbb{P}}(\pi, \varphi) \leq P_{\mathbb{P}}(\pi, \mathbf{0}) + \max \varphi$.
- (v) $P_{\mathbb{P}}(\pi, \cdot)$ is either finite valued or constantly $\pm\infty$.
- (vi) If $P_{\mathbb{P}}(\pi, \cdot) \neq \pm\infty$, then $|P_{\mathbb{P}}(\pi, \varphi) - P_{\mathbb{P}}(\pi, \psi)| \leq \|\varphi - \psi\|$.
- (vii) If $P_{\mathbb{P}}(\pi, \cdot) \neq \pm\infty$, then $P_{\mathbb{P}}(\pi, \cdot)$ is convex.
- (viii) $P_{\mathbb{P}}(\pi, c\varphi) \leq c \cdot P_{\mathbb{P}}(\pi, \varphi)$, if $c \geq 1$ and $P_{\mathbb{P}}(\pi, c\varphi) \geq c \cdot P_{\mathbb{P}}(\pi, \varphi)$, if $c \leq 1$.
- (ix) $|P_{\mathbb{P}}(\pi, \varphi)| \leq P_{\mathbb{P}}(\pi, |\varphi|)$.
- (x) $P_{\mathbb{P}}(\pi, \psi + \varphi \circ f - \varphi) = P_{\mathbb{P}}(\pi, \psi)$, for any $f \in [[\mathcal{H}]]_{top}$.

Acknowledgements. The work was supported by the National Natural Science Foundation of China (grant No. 11271191) and National Basic Research Program of China (grant No. 2013CB834100) and the Foundation for Innovative program of Jiangsu province (grant No. CXZZ12 0380).

References

- [1] L. Bowen, Measure conjugacy invariants for actions of countable sofic groups, J. Amer. Math. Soc., **23** (2010), no. 1, 217-245.
- [2] L. Bowen, Sofic entropy and amenable groups, Ergod. Th. Dynam. Sys., to appear.
- [3] L. Bowen, Entropy theory for sofic groupoids I: the foundations, preprint.
- [4] R. Bowen, Topological entropy for non-compact set, Trans. Amer. Math. Soc., **184** (1973), 125-136.
- [5] N. Chung, The variational principle of topological pressure for actions of sofic groups. arXiv: 1110.0699v1.
- [6] E. Glasner & B. Weiss, On the interplay between measurable and topological dynamics, in Handbook of dynamical systems, Vol. 1B, Elsevier B. V., Amsterdam, (2006) 597-648.
- [7] T. Goodman, Topological entropy bounds measure-theoretic entropy, Proc. Amer. Math. Soc., **23** (1969), 679-688.
- [8] L. Goodwyn, Relating topological entropy and measure entropy, Bull. London Math. Soc., **3** (1971), 176-180.
- [9] W. Huang & X. Ye, A local variational relation and applications, Israel J. Math., **151** (2006), 237-279.
- [10] W. Huang, X. Ye & G. Zhang, Local entropy theory for a countable discrete amenable group action, J. Funct. Anal., **261**(2011), 1028-1082.
- [11] W. Huang & Y. Yi, A Local variational principle of pressure and its applications to equilibrium states, Israel J. Math., **161** (2007), 29-74.
- [12] D. Kerr, Sofic measure entropy via finite partitions, Geometry, Group and Dynamics, to appear.
- [13] D. Kerr & H. Li, Entropy and the variational principle for actions of sofic groups, Invent. Math., **186** (2011), 501–558.
- [14] D. Kerr & H. Li, Soficity, amenability, and dynamical entropy, Amer. J. Math., to appear.
- [15] H. Li, Sofic mean dimension, arXiv: 1105.0140v1.

- [16] B. Liang & K. Yan, Topological pressure for sub-additive potentials of amenable group actions, *J. Func. Anal.*, **262** (2012), 584-601.
- [17] M. Misiurewicz, A short proof of the variational principle for a $\mathbb{Z}_+^n$ action on a compact space, *Bull. Acad. Polon. Sci. Sr. Sci. Math. Astronom. Phys.*, **24** (1976), 1069-1075.
- [18] D. Ornstein & B. Weiss, Entropy and isomorphism theorems for actions of amenable groups, *J. Anal. Math.*, **48** (1987), 1-141.
- [19] Y. Pesin & B. Pitskel, Topological pressure and the variational principle for non-compact sets, *Func. Anal. Appl.*, **18** (1984), 307-318.
- [20] P. Romagnoli, A local variational principle for the topological entropy, *Ergod. Th. Dynam. Sys.*, **23** (2003), 1601-1610.
- [21] P. Walter, A variational principle for the pressure of continuous transformations, *Amer. J. Math.*, **97** (1975), 937-971.
- [22] P. Walters, *An Introduction to Ergodic Theory*, Springer, New York, 1982.
- [23] G. Zhang, Local variational principle concerning entropy of a sofic group action, *J. Func. Anal.*, **262** (2012), 1954-1985.